

Products of Many Sorted Algebras

Beata Madras
Warsaw University
Białystok

Summary. Product of two many sorted universal algebras and product of family of many sorted universal algebras are defined in this article. Operations on functions, such that commute, Frege, are also introduced.

MML Identifier: PRALG_2.

WWW: http://mizar.org/JFM/Vol6/pralg_2.html

The articles [13], [9], [19], [6], [7], [14], [18], [10], [8], [15], [4], [11], [1], [3], [2], [5], [16], [12], and [17] provide the notation and terminology for this paper.

1. PRELIMINARIES

For simplicity, we follow the rules: I, J are sets, A, B are many sorted sets indexed by I , i, j, x are sets, and S is a non empty many sorted signature.

Let I_1 be a set. We say that I_1 has common domain if and only if:

(Def. 1) For all functions f, g such that $f \in I_1$ and $g \in I_1$ holds $\text{dom } f = \text{dom } g$.

Let us note that there exists a set which is functional and non empty and has common domain.

The following proposition is true

(1) $\{\emptyset\}$ is a functional set with common domain.

Let X be a functional set with common domain. The functor $\text{DOM}(X)$ yielding a set is defined by:

(Def. 2)(i) For every function x such that $x \in X$ holds $\text{DOM}(X) = \text{dom } x$ if $X \neq \emptyset$,

(ii) $\text{DOM}(X) = \emptyset$, otherwise.

Let X be a functional non empty set with common domain. We see that the element of X is a many sorted set indexed by $\text{DOM}(X)$.

One can prove the following proposition

(2) For every functional set X with common domain such that $X = \{\emptyset\}$ holds $\text{DOM}(X) = \emptyset$.

Let I be a set and let M be a non-empty many sorted set indexed by I . One can verify that $\prod M$ is functional and non empty and has common domain.

2. OPERATIONS ON FUNCTIONS

The scheme *LambdaDMS* deals with a non empty set \mathcal{A} and a unary functor \mathcal{F} yielding a set, and states that:

There exists a many sorted set X indexed by \mathcal{A} such that for every element d of \mathcal{A} holds $X(d) = \mathcal{F}(d)$

for all values of the parameters.

Let f be a function. The functor $\text{commute}(f)$ yielding a function yielding function is defined by:

(Def. 5)¹ $\text{commute}(f) = \text{curry}' \text{uncurry } f$.

The following propositions are true:

- (3) For every function f and for every set x such that $x \in \text{dom } \text{commute}(f)$ holds $(\text{commute}(f))(x)$ is a function.
- (4) For all sets A, B, C and for every function f such that $A \neq \emptyset$ and $B \neq \emptyset$ and $f \in (C^B)^A$ holds $\text{commute}(f) \in (C^A)^B$.
- (5) Let A, B, C be sets and f be a function. Suppose $A \neq \emptyset$ and $B \neq \emptyset$ and $f \in (C^B)^A$. Let g, h be functions and x, y be sets. Suppose $x \in A$ and $y \in B$ and $f(x) = g$ and $(\text{commute}(f))(y) = h$. Then $h(x) = g(y)$ and $\text{dom } h = A$ and $\text{dom } g = B$ and $\text{rng } h \subseteq C$ and $\text{rng } g \subseteq C$.
- (6) For all sets A, B, C and for every function f such that $A \neq \emptyset$ and $B \neq \emptyset$ and $f \in (C^B)^A$ holds $\text{commute}(\text{commute}(f)) = f$.
- (7) $\text{commute}(\square) = \square$.

Let F be a function. The functor $\blacksquare \text{commute}(F)$ yielding a function is defined by the conditions (Def. 6).

- (Def. 6)(i) For every x holds $x \in \text{dom } \blacksquare \text{commute}(F)$ iff there exists a function f such that $f \in \text{dom } F$ and $x = \text{commute}(f)$, and
- (ii) for every function f such that $f \in \text{dom } \blacksquare \text{commute}(F)$ holds $(\blacksquare \text{commute}(F))(f) = F(\text{commute}(f))$.

The following proposition is true

- (8) For every function F such that $\text{dom } F = \{\emptyset\}$ holds $\blacksquare \text{commute}(F) = F$.

Let F be a function yielding function and let f be a function. The functor $F \leftarrow \wp f$ yields a function and is defined as follows:

- (Def. 7) $\text{dom}(F \leftarrow \wp f) = \text{dom } F$ and for every set x such that $x \in \text{dom } F$ holds $(F \leftarrow \wp f)(x) = F(x)(f(x))$.

Let f be a function yielding function. Then $\text{Frege}(f)$ is a many sorted function indexed by $\prod(\text{dom}_\kappa f(\kappa))$ and it can be characterized by the condition:

- (Def. 8) For every function g such that $g \in \prod(\text{dom}_\kappa f(\kappa))$ holds $(\text{Frege}(f))(g) = f \leftarrow \wp g$.

Let us consider I, A, B . The functor $\llbracket A, B \rrbracket$ yields a many sorted set indexed by I and is defined by:

- (Def. 9) For every i such that $i \in I$ holds $\llbracket A, B \rrbracket(i) = \{ A(i), B(i) \}$.

Let us consider I and let A, B be non-empty many sorted sets indexed by I . Observe that $\llbracket A, B \rrbracket$ is non-empty.

Next we state the proposition

¹ The definitions (Def. 3) and (Def. 4) have been removed.

- (9) Let I be a non empty set, J be a set, A, B be many sorted sets indexed by I , and f be a function from J into I . Then $\llbracket A, B \rrbracket \cdot f = \llbracket A \cdot f, B \cdot f \rrbracket$.

Let I be a non empty set, let us consider J , let A, B be non-empty many sorted sets indexed by I , let p be a function from J into I^* , let r be a function from J into I , let j be a set, let f be a function from $(A^\# \cdot p)(j)$ into $(A \cdot r)(j)$, and let g be a function from $(B^\# \cdot p)(j)$ into $(B \cdot r)(j)$. Let us assume that $j \in J$. The functor $\llbracket f, g \rrbracket$ yielding a function from $(\llbracket A, B \rrbracket^\# \cdot p)(j)$ into $(\llbracket A, B \rrbracket \cdot r)(j)$ is defined by:

- (Def. 10) For every function h such that $h \in ((\llbracket A, B \rrbracket^\# \cdot p)(j))$ holds $\llbracket f, g \rrbracket(h) = \langle f(\text{pr1}(h)), g(\text{pr2}(h)) \rangle$.

Let I be a non empty set, let us consider J , let A, B be non-empty many sorted sets indexed by I , let p be a function from J into I^* , let r be a function from J into I , let F be a many sorted function from $A^\# \cdot p$ into $A \cdot r$, and let G be a many sorted function from $B^\# \cdot p$ into $B \cdot r$. The functor $\llbracket F, G \rrbracket$ yields a many sorted function from $\llbracket A, B \rrbracket^\# \cdot p$ into $\llbracket A, B \rrbracket \cdot r$ and is defined by the condition (Def. 11).

- (Def. 11) Let given j . Suppose $j \in J$. Let f be a function from $(A^\# \cdot p)(j)$ into $(A \cdot r)(j)$ and g be a function from $(B^\# \cdot p)(j)$ into $(B \cdot r)(j)$. If $f = F(j)$ and $g = G(j)$, then $\llbracket F, G \rrbracket(j) = \llbracket f, g \rrbracket$.

3. FAMILY OF MANY SORTED UNIVERSAL ALGEBRAS

Let us consider I, S . A many sorted set indexed by I is said to be an algebra family of I over S if:

- (Def. 12) For every i such that $i \in I$ holds $\text{it}(i)$ is a non-empty algebra over S .

Let I be a non empty set, let us consider S , let A be an algebra family of I over S , and let i be an element of I . Then $A(i)$ is a non-empty algebra over S .

Let S be a non empty many sorted signature and let U_1 be an algebra over S . The functor $|U_1|$ yielding a set is defined by:

- (Def. 13) $|U_1| = \bigcup \text{rng}(\text{the sorts of } U_1)$.

Let S be a non empty many sorted signature and let U_1 be a non-empty algebra over S . Observe that $|U_1|$ is non empty.

Let I be a non empty set, let S be a non empty many sorted signature, and let A be an algebra family of I over S . The functor $|A|$ yielding a set is defined as follows:

- (Def. 14) $|A| = \bigcup \{ |A(i)| : i \text{ ranges over elements of } I \}$.

Let I be a non empty set, let S be a non empty many sorted signature, and let A be an algebra family of I over S . Observe that $|A|$ is non empty.

4. PRODUCT OF MANY SORTED UNIVERSAL ALGEBRAS

Next we state two propositions:

- (10) Let S be a non void non empty many sorted signature, U_0 be an algebra over S , and o be an operation symbol of S . Then $\text{Args}(o, U_0) = \prod(\text{the sorts of } U_0 \cdot \text{Arity}(o))$ and $\text{dom}(\text{the sorts of } U_0 \cdot \text{Arity}(o)) = \text{dom Arity}(o)$ and $\text{Result}(o, U_0) = (\text{the sorts of } U_0)(\text{the result sort of } o)$.

- (11) Let S be a non void non empty many sorted signature, U_0 be an algebra over S , and o be an operation symbol of S . If $\text{Arity}(o) = \varepsilon$, then $\text{Args}(o, U_0) = \{\square\}$.

Let us consider S and let U_1, U_2 be non-empty algebras over S . The functor $\{U_1, U_2\}$ yielding an algebra over S is defined as follows:

(Def. 15) $\{U_1, U_2\} = \langle \llbracket \text{the sorts of } U_1, \text{ the sorts of } U_2 \rrbracket, \llbracket \text{the characteristics of } U_1, \text{ the characteristics of } U_2 \rrbracket \rangle$.

Let us consider S and let U_1, U_2 be non-empty algebras over S . One can verify that $\{U_1, U_2\}$ is strict.

Let us consider I, S , let s be a sort symbol of S , and let A be an algebra family of I over S . The functor $\text{Carrier}(A, s)$ yielding a many sorted set indexed by I is defined as follows:

- (Def. 16)(i) For every set i such that $i \in I$ there exists an algebra U_0 over S such that $U_0 = A(i)$ and $(\text{Carrier}(A, s))(i) = (\text{the sorts of } U_0)(s)$ if $I \neq \emptyset$,
(ii) $\text{Carrier}(A, s) = \emptyset$, otherwise.

Let us consider I, S , let s be a sort symbol of S , and let A be an algebra family of I over S . Note that $\text{Carrier}(A, s)$ is non-empty.

Let us consider I, S and let A be an algebra family of I over S . The functor $\text{SORTS}(A)$ yields a many sorted set indexed by the carrier of S and is defined as follows:

(Def. 17) For every sort symbol s of S holds $(\text{SORTS}(A))(s) = \coprod \text{Carrier}(A, s)$.

Let us consider I, S and let A be an algebra family of I over S . Observe that $\text{SORTS}(A)$ is non-empty.

Let us consider I , let S be a non empty many sorted signature, and let A be an algebra family of I over S . The functor $\text{OPER}(A)$ yielding a many sorted function indexed by I is defined as follows:

- (Def. 18)(i) For every set i such that $i \in I$ there exists an algebra U_0 over S such that $U_0 = A(i)$ and $(\text{OPER}(A))(i) = \text{the characteristics of } U_0$ if $I \neq \emptyset$,
(ii) $\text{OPER}(A) = \emptyset$, otherwise.

We now state two propositions:

- (12) Let S be a non empty many sorted signature and A be an algebra family of I over S . Then $\text{dom uncurry OPER}(A) = \{I, \text{the operation symbols of } S\}$.
(13) Let I be a non empty set, S be a non void non empty many sorted signature, A be an algebra family of I over S , and o be an operation symbol of S . Then $\text{commute}(\text{OPER}(A)) \in ((\text{rng uncurry OPER}(A))^I)^{\text{the operation symbols of } S}$.

Let I be a set, let S be a non void non empty many sorted signature, let A be an algebra family of I over S , and let o be an operation symbol of S . The functor $A(o)$ yields a many sorted function indexed by I and is defined as follows:

(Def. 19) $A(o) = (\text{commute}(\text{OPER}(A)))(o)$.

The following propositions are true:

- (14) Let I be a non empty set, i be an element of I , S be a non void non empty many sorted signature, A be an algebra family of I over S , and o be an operation symbol of S . Then $A(o)(i) = \text{Den}(o, A(i))$.
(15) Let I be a non empty set, S be a non void non empty many sorted signature, A be an algebra family of I over S , o be an operation symbol of S , and x be a set. If $x \in \text{rng Frege}(A(o))$, then x is a function.
(16) Let I be a non empty set, S be a non void non empty many sorted signature, A be an algebra family of I over S , o be an operation symbol of S , and f be a function. If $f \in \text{rng Frege}(A(o))$, then $\text{dom } f = I$ and for every element i of I holds $f(i) \in \text{Result}(o, A(i))$.

- (17) Let I be a non empty set, S be a non void non empty many sorted signature, A be an algebra family of I over S , o be an operation symbol of S , and f be a function. Suppose $f \in \text{dom Frege}(A(o))$. Then $\text{dom } f = I$ and for every element i of I holds $f(i) \in \text{Args}(o, A(i))$ and $\text{rng } f \subseteq |A|^{\text{dom Arity}(o)}$.
- (18) Let I be a non empty set, S be a non void non empty many sorted signature, A be an algebra family of I over S , and o be an operation symbol of S . Then $\text{dom}(\text{dom}_\kappa A(o)(\kappa)) = I$ and for every element i of I holds $(\text{dom}_\kappa A(o)(\kappa))(i) = \text{Args}(o, A(i))$.

Let us consider I , let S be a non void non empty many sorted signature, and let A be an algebra family of I over S . The functor $\text{OPS}(A)$ yielding a many sorted function from $(\text{SORTS}(A))^\#$ · the arity of S into $\text{SORTS}(A)$ · the result sort of S is defined by:

- (Def. 20)(i) For every operation symbol o of S holds $(\text{OPS}(A))(o) = (\text{Arity}(o) = \varepsilon \rightarrow \text{commute}(A(o)), \blacksquare \text{commute}(\text{Frege}(A(o))))$ if $I \neq \emptyset$,
- (ii) TRUE, otherwise.

Let I be a set, let S be a non void non empty many sorted signature, and let A be an algebra family of I over S . The functor $\prod A$ yielding an algebra over S is defined as follows:

- (Def. 21) $\prod A = \langle \text{SORTS}(A), \text{OPS}(A) \rangle$.

Let I be a set, let S be a non void non empty many sorted signature, and let A be an algebra family of I over S . Note that $\prod A$ is strict.

The following propositions are true:

- (19) For every non void non empty many sorted signature S and for every algebra family A of I over S holds $\prod A = \langle \text{SORTS}(A), \text{OPS}(A) \rangle$.
- (20) Let S be a non void non empty many sorted signature and A be an algebra family of I over S . Then the sorts of $\prod A = \text{SORTS}(A)$ and the characteristics of $\prod A = \text{OPS}(A)$.

REFERENCES

- [1] Grzegorz Bancerek. Curried and uncurried functions. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_5.html.
- [2] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [3] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/funct_6.html.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [5] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/dtconstr.html>.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [7] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [8] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [9] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [10] Czesław Byliński. A classical first order language. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/cqc_lang.html.
- [11] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.

- [12] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [14] Andrzej Trybulec. Tuples, projections and Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/mcart_1.html.
- [15] Andrzej Trybulec. Function domains and Frænkel operator. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/fraenkel.html>.
- [16] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pbool1e.html>.
- [17] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [18] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [19] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received April 25, 1994

Published July 11, 2000
