

THREE KINDS OF MUTATION

ASLAK BAKKE BUAN, IDUN REITEN, AND HUGH THOMAS

ABSTRACT. For a finite dimensional hereditary algebra H , we consider: exceptional sequences in $\text{mod } H$, silting objects in the bounded derived category \mathcal{D} , and m -cluster tilting objects in the m -cluster category, $\mathcal{C}_m = \mathcal{D}/\tau^{-1}[m]$. There are mutation operations on both the set of m -cluster tilting objects and the set of exceptional sequences. It is also possible to define a mutation operation for silting objects. We compare these three different notions of mutation.

INTRODUCTION

Let H be a hereditary finite dimensional algebra with n isomorphism classes of simple objects. The tilting objects for H inside $\mathcal{D} = D^b(H)$, the bounded derived category of finitely generated H modules, are of considerable importance, and several variations on the concept have also been studied. In this paper, we will be concerned with three of these: exceptional sequences in $\text{mod } H$, silting objects in \mathcal{D} , and m -cluster tilting objects in the m -cluster category, $\mathcal{C}_m = \mathcal{D}/\tau^{-1}[m]$, which is an orbit category of \mathcal{D} .

There are mutation operations on both the set of m -cluster tilting objects and the set of exceptional sequences. It is also possible to define a mutation operation for silting objects. Our main goal in this paper is to relate these different notions of mutation. In order to state our results, we must introduce some definitions.

A basic object T in \mathcal{D} is said to be partial silting if $\text{Ext}^i(T, T) = 0$ for $i > 0$, and silting if in addition T is maximal with this property. Partial silting objects were also studied in [KV] (they were then called silting objects), and in [AST], where some connections to exceptional sequences were also investigated.

A basic object T in \mathcal{C}_m is called m -cluster tilting if $\text{Ext}_{\mathcal{C}_m}^i(T, T) = 0$ for $0 < i \leq m$ and if X satisfies $\text{Ext}_{\mathcal{C}_m}^i(T, X) = 0$ for $0 < i \leq m$, then X is in $\text{add } T$. This condition is known to be equivalent to T being maximal with respect to $\text{Ext}_{\mathcal{C}_m}^i(T, T) = 0$ for $0 < i \leq m$ [W, ZZ], and to T having n indecomposable summands. If T satisfies $\text{Ext}_{\mathcal{C}_m}^i(T, T) = 0$ for $0 < i \leq m$ and it has $n - 1$ summands, it is called *almost m -cluster tilting*.

If we make a (reasonable, but not canonical) choice of a fundamental domain in \mathcal{D} for $\tau^{-1}[m]$, then the lifting of objects from \mathcal{C}_m to \mathcal{D} takes m -cluster tilting objects to silting objects. Conversely, any silting object is the lift of an m -cluster tilting object for m sufficiently large. This allows us to deduce the number of summands of a silting object, and to lift the notion of mutation from m -cluster tilting objects to silting objects.

Date: May 4, 2010.

Exceptional sequences were investigated first in the context of algebraic geometry (see [Ru]), and later for hereditary finite dimensional algebras in [CB, R]. An exceptional sequence in $\text{mod } H$ is a sequence of indecomposable H modules, (E_1, \dots, E_r) , with the properties that $\text{Hom}(E_i, E_j) = 0$ for $j < i$ and $\text{Ext}^1(E_i, E_j) = 0$ for $j \leq i$. The maximal length of an exceptional sequence is n ; an exceptional sequence of maximal length is called *complete*, and one of length $n - 1$ is called *almost complete*.

Given a sequence of indecomposable H modules (M_1, \dots, M_n) , a *placement* of (M_1, \dots, M_n) is a sequence $(\widehat{M}_1, \dots, \widehat{M}_n)$ of objects in \mathcal{D} such that \widehat{M}_i is isomorphic to $M_i[t_i]$ for some $t_i \in \mathbb{Z}$. We will be interested in studying placements of an exceptional sequence which form a silting object. It is easy to see that any complete exceptional sequence admits such a placement. It is natural to ask if it is possible to define a placement for each exceptional sequence in $\text{mod } H$, such that if two exceptional sequences are related by a single mutation, the same would be true for the corresponding silting objects. Simple examples show that this is too much to ask. In fact, for some H , there exists an exceptional sequence \mathcal{E} such that it is not possible to place \mathcal{E} and the sequences \mathcal{E}_i obtained by applying a single mutation to \mathcal{E} , in such a way that the corresponding silting object $\widehat{\mathcal{E}}$ is related by a single mutation to each of the $\widehat{\mathcal{E}}_i$.

Our main result is as follows. Let \mathcal{F} be an almost complete exceptional sequence, and $\mathcal{E}_1, \dots, \mathcal{E}_{n+1}$ be the complete exceptional sequences which can be obtained by adding one term to \mathcal{F} . Essentially by definition, \mathcal{E}_i and \mathcal{E}_{i+1} are related by a single mutation. Then there exists a placement $\widehat{\mathcal{E}}_i$ of each \mathcal{E}_i , all of which agree as to the placement of \mathcal{F} , such that $\widehat{\mathcal{E}}_i$ is silting for all i , and such that $\widehat{\mathcal{E}}_i$ and $\widehat{\mathcal{E}}_{i+1}$ are related by a single mutation.

In the course of our investigations, we define a quiver associated to an exceptional sequence, as follows. The vertex set consists of the terms in the exceptional sequence; there is an arrow from E_i to E_j if $\text{Hom}(E_i, E_j) \neq 0$ or $\text{Ext}^1(E_j, E_i) \neq 0$. (Note the reversal of the order of E_i and E_j in the two conditions.) We show that this quiver is always acyclic, and that it is connected provided H is connected and the exceptional sequence is complete. This acyclicity is crucial for our main result mentioned above.

In section 1, we discuss exceptional sequences, starting with definitions and basic properties and we define and study the quiver associated with an exceptional sequence. In section 2 we give connections between exceptional sequences, silting objects and m -cluster tilting objects and in section 3 we consider connections between the different kinds of mutation. Finally, in section 4 we show that for a fixed almost complete exceptional sequence \mathcal{E} , and its set of complements, there is a natural interpretation of \mathcal{E} as an almost complete silting object in such a way that complements and exchange sequences correspond.

1. PROPERTIES OF EXCEPTIONAL SEQUENCES

In this section we first recall the definition and some basic properties of exceptional sequences. Then we define, for each exceptional sequence \mathcal{E} , a naturally associated quiver $\mathcal{Q}_{\mathcal{E}}$, and show that such quivers are always acyclic, and are connected provided

H is connected and \mathcal{E} is complete. In section 4, these quivers and their acyclicity will be a main ingredient in the proof of Theorem 4.1.

As before, let H be a finite dimensional hereditary algebra over a field k . We do not assume that k is algebraically closed. We always assume that H has n simple modules up to isomorphism. An indecomposable module E is called *exceptional* if $\text{Ext}^1(E, E) = 0$, and a sequence (E_1, \dots, E_r) of exceptional H -modules is said to be an *exceptional sequence* if $\text{Hom}(E_j, E_i) = 0 = \text{Ext}^1(E_j, E_i)$ for $j > i$.

1.1. Basic properties. We have the following useful fact.

Lemma 1.1. *Given an exceptional sequence (E_1, \dots, E_r) , then $\text{Hom}(E_i, E_j)$ and $\text{Ext}^1(E_i, E_j)$ cannot both be non-zero for $i < j$.*

Proof. Fix $E = E_i$, $F = E_j$. By results from [CB, R], we can consider (E, F) as an exceptional sequence in a hereditary module category, say $\text{mod } H'$, with H' of rank 2 and such that $\text{mod } H'$ has a full and exact embedding into $\text{mod } H$. For a hereditary algebra H' of rank 2, the only exceptional indecomposable modules are preprojective or preinjective. Hence, a case analysis of the possible exceptional sequences in $\text{mod } H'$ for such algebras, gives the result. \square

There are right and left mutation operations on exceptional sequences. Here we use *right mutation* μ_i , which for an exceptional sequence $\mathcal{E} = (E_1, E_2, \dots, E_r)$ and an index $i \in \{1, \dots, r-1\}$ is defined by replacing the pair (E_i, E_{i+1}) with a pair (E_{i+1}, E_i^*) . Here E_i^* is defined as follows. If $\text{Hom}(E_i, E_{i+1}) \neq 0$, consider the minimal left add E_{i+1} -approximation $f: E_i \rightarrow E_{i+1}^t$. Then f is either an epimorphism or a monomorphism (by [HR, RS]). If f is an epimorphism, let $E_i^* = \ker f$. If f is a monomorphism, let $E_i^* = \text{coker } f$. If $\text{Ext}^1(E_i, E_{i+1}) \neq 0$, then E_i^* is defined by the universal extension [B]

$$0 \rightarrow E_{i+1}^s \rightarrow E_i^* \rightarrow E_i \rightarrow 0.$$

Finally, if $\text{Hom}(E_i, E_{i+1}) = 0 = \text{Ext}^1(E_i, E_{i+1})$, then $E_i^* = E_i$.

One obtains a uniform description of E_i^* as follows. Let thick E_{i+1} be the thick subcategory of \mathcal{D} generated by E_{i+1} . The indecomposable objects in this subcategory are of the form $E_{i+1}[j]$, for some integer j . Let $E_i \rightarrow Z$ be a minimal left (thick E_{i+1})-approximation in \mathcal{D} , and complete to a triangle

$$E_i \rightarrow Z \rightarrow E_i^\sharp.$$

It is easy to see that $E_i^\sharp = E_i^*[j]$ with j either 0 or 1.

We recall the following from [CB, R].

Proposition 1.2. *Let $\mathcal{E} = (E_1, E_2, \dots, E_r)$ be an exceptional sequence in $\text{mod } H$.*

- (a) *We have $r \leq n$.*
- (b) *If $r < n$, then there is an complete exceptional sequence*

$$(E_1, E_2, \dots, E_r, E_{r+1}, \dots, E_n).$$

- (c) *If $r = n - 1$, then there is for each $j \in \{1, \dots, n\}$ a unique indecomposable module M , such that*

$$(E_1, \dots, E_{j-1}, M, E_j, \dots, E_{n-1})$$

is an exceptional sequence.

- (d) There is an ordering S_1, \dots, S_n of the simple H -modules, such that $\mathcal{S} = (S_1, \dots, S_n)$ is a complete exceptional sequence. Moreover, any complete exceptional sequence can be reached from \mathcal{S} by a sequence of mutations and inverses of mutations.

1.2. The Hom-Ext-quiver. We now associate a quiver $\mathcal{Q}_{\mathcal{E}}$ to an exceptional sequence $\mathcal{E} = (E_1, E_2, \dots, E_r)$. The vertices of $\mathcal{Q}_{\mathcal{E}}$ are in bijection with $\{E_1, E_2, \dots, E_r\}$. There is an arrow $E_i \rightarrow E_j$ if $\text{Hom}(E_i, E_j) \neq 0$ or if $\text{Ext}^1(E_j, E_i) \neq 0$. We call this the *Hom-Ext quiver* of \mathcal{E} .

The construction of this quiver was motivated by the situation when H is representation finite. In that case, the AR quiver defines a partial order on the indecomposable H -modules, which can be restricted to give a partial order on the terms of \mathcal{E} . If $E_i \rightarrow E_j$ in the Hom-Ext quiver, then E_i precedes E_j with respect to this partial order (though not conversely). Where we have used the acyclicity of $\mathcal{Q}_{\mathcal{E}}$ below, we could instead have used the order induced from the AR quiver. When H is not representation finite, however, the AR quiver will have more than one component and will typically contain cycles, making it difficult to use it to define a partial order on the terms of \mathcal{E} ; the Hom-Ext quiver is a replacement for this order.

Note that if A and B are distinct exceptional modules with $\text{Hom}(A, B) \neq 0$, then there is, by [HR], either a monomorphism $A \rightarrow B$ or an epimorphism $A \rightarrow B$ and (by reason of total dimension) not both. Let us add some decoration to our quiver. There are three types of arrows. There is an *m*-arrow $E_i \rightarrow E_j$ if there is a monomorphism $E_i \rightarrow E_j$. There is an *e*-arrow $E_i \rightarrow E_j$ if there is an epimorphism $E_i \rightarrow E_j$, and there is an *x*-arrow $E_i \rightarrow E_j$ if $\text{Ext}^1(E_j, E_i) \neq 0$.

We aim to prove that the quiver $\mathcal{Q}_{\mathcal{E}}$ is acyclic and that it is connected provided \mathcal{E} is complete and H is connected. For this purpose the following result is useful.

Lemma 1.3. *Assume A, B, C are exceptional modules contained in an exceptional sequence \mathcal{E} . Then we have the following.*

- (a) *If there is an e-arrow $A \rightarrow B$, then there is not an m-arrow $B \rightarrow C$.*
- (b) *If there are m-arrows $A \rightarrow B$ and $B \rightarrow C$, there must be an m-arrow $A \rightarrow C$.*
- (c) *If there are e-arrows $A \rightarrow B$ and $B \rightarrow C$, there must be an e-arrow $A \rightarrow C$.*
- (d) *If there is an e-arrow $A \rightarrow B$ and an x-arrow $B \rightarrow C$, then there is an x-arrow $A \rightarrow C$.*
- (e) *If there is an x-arrow $A \rightarrow B$ and an m-arrow $B \rightarrow C$, then there is an x-arrow $A \rightarrow C$.*
- (f) *If there are arrows $A \rightarrow B \rightarrow C$ and $B \rightarrow C$ is an m-arrow, then there is an arrow $A \rightarrow C$.*

Proof. (a) follows from the fact that the composition of an epimorphism and a monomorphism is a nonzero map which is neither a monomorphism nor an epimorphism.

(b) and (c) are obvious.

For (d) apply $\text{Hom}(C, _)$ to the exact sequence

$$0 \rightarrow K \rightarrow A \rightarrow B \rightarrow 0$$

to obtain an epimorphism $\text{Ext}^1(C, A) \rightarrow \text{Ext}^1(C, B)$.

(e) is dual to (d).

For (f), we have that $A \rightarrow B$ cannot be an e-arrow by (a). Then the claim follows from combining (b) and (e). \square

Applying this we first show acyclicity.

Theorem 1.4. *The Hom-Ext quiver $\mathcal{Q}_{\mathcal{E}}$ of a complete exceptional sequence is acyclic.*

Proof. Assume there is a cycle in the quiver $\mathcal{Q}_{\mathcal{E}}$. We can assume that the cycle has minimal length. Hence, for any two vertices X and Y on this cycle which are not consecutive on the cycle, there are no arrows between X and Y . Note also that by Lemma 1.1 the length must be at least three. By Lemma 1.3 (f) there cannot be any m-arrows on the minimal cycle.

It is clear that there must be at least one x-arrow on the cycle, and at least one arrow which is not an x-arrow, since otherwise the objects corresponding to vertices on the cycles could not be ordered into an exceptional sequence. Thus there must be an x-arrow which is preceded by an e-arrow. However, by Lemma 1.3 (e), this is not possible on a minimal cycle. \square

We also have the following property.

Theorem 1.5. *If H is connected, then the Hom-Ext quiver $\mathcal{Q}_{\mathcal{E}}$ of a complete exceptional sequence \mathcal{E} in $\text{mod } H$ is connected.*

Proof. Assume that $\mathcal{Q}_{\mathcal{E}}$ is not connected, and assume that $\mathcal{E} = \mathcal{E}_1 \cup \mathcal{E}_2$, with $\mathcal{E}_1 = \{A_1, \dots, A_r\}$ and $\mathcal{E}_2 = \{B_1, \dots, B_s\}$ being non-empty disjoint subsets of the elements in \mathcal{E} , with the property that there are no arrows between any vertex corresponding to an object in \mathcal{E}_1 and any vertex corresponding to an object in \mathcal{E}_2 .

This is equivalent to having $\text{Hom}(A_i, B_j) = \text{Hom}(B_j, A_i) = \text{Ext}^1(A_i, B_j) = \text{Ext}^1(B_j, A_i) = 0$ for all i, j . We claim that any exceptional sequence \mathcal{E}' obtained by mutating \mathcal{E} , also has the property that $\mathcal{Q}_{\mathcal{E}'}$ is disconnected. This will give a contradiction, because of transitivity of the action of mutations on exceptional sequences, and Proposition 1.2 (d). To prove the claim consider the mutation pair (E_i, E_{i+1}) , and consider the four possibilities:

- (1) $\text{Hom}(E_i, E_{i+1}) = \text{Ext}^1(E_i, E_{i+1}) = 0$ and $E_i^* = E_i$
- (2) There is an exact sequence $0 \rightarrow E_i \rightarrow E_{i+1}^t \rightarrow E_i^* \rightarrow 0$
- (3) There is an exact sequence $0 \rightarrow E_i^* \rightarrow E_i \rightarrow E_{i+1}^t \rightarrow 0$
- (4) There is an exact sequence $0 \rightarrow E_{i+1}^t \rightarrow E_i^* \rightarrow E_i \rightarrow 0$

Recall that in cases (2) and (3) the map $E_i \rightarrow E_{i+1}^t$ is a minimal left add E_{i+1} -approximation, and in case (4) the exact sequence is a universal extension.

We claim that if we replace E_i with E_i^* , then $\mathcal{Q}_{\mathcal{E}'}$ will disconnect in the same way as $\mathcal{Q}_{\mathcal{E}}$. If we are in situation (1) above, this is trivial. Assume we are in situation (2), (3) or (4). Then either both E_i and E_{i+1} are in \mathcal{E}_1 or both are in \mathcal{E}_2 . Without loss of generality we assume they are both in \mathcal{E}_1 . Let B an object in \mathcal{E}_2 . Considering the long exact sequences obtained from applying $\text{Hom}(B, _)$ and $\text{Hom}(_, B)$ to the relevant exact sequence (2), (3) or (4), we obtain that $\text{Hom}(E_i^*, B) = \text{Hom}(B, E_i^*) = \text{Ext}^1(E_i^*, B) = \text{Ext}^1(B, E_i^*) = 0$. This gives the desired result.

□

2. CONNECTION BETWEEN EXCEPTIONAL SEQUENCES, SILTING OBJECTS AND m -CLUSTER TILTING OBJECTS

In this section we point out how to construct silting objects from exceptional sequences and vice versa. We also investigate the connection between silting objects and m -cluster tilting objects.

2.1. Silting objects and exceptional sequences. The results in this section partially overlap with [AST]. As before, let H be a finite dimensional hereditary algebra with n isomorphism classes of simples, and let T be an object in \mathcal{D} . Recall that T is called a *partial silting object* if $\text{Ext}_{\mathcal{D}}^i(T, T) = 0$ for all $i > 0$. Without loss of generality we can assume that T is in the non-negative part \mathcal{D}^+ of \mathcal{D} , i.e. we assume

$$(1) \quad T = T_0[0] \oplus T_1[1] \oplus \cdots \oplus T_m[m]$$

with each T_i in $\text{mod } H$. Then we have the following.

Lemma 2.1. *T is partial silting if and only if*

- (i) $\text{Ext}_H^1(T_i, T_i) = 0$ for each i
- (ii) $\text{Hom}(T_i, T_j) = 0 = \text{Ext}_H^1(T_i, T_j)$ for $i > j$.

Proof. We have that T is partial silting if and only if $\text{Ext}^1(T_i, T_i) = 0$ for each i and $\text{Hom}(T_i[i], T_j[j][t]) = \text{Hom}(T_i[i], T_j[t+j]) = 0$ for $t > 0$. Note that if $i < j$, then $\text{Hom}(T_i[i], T_j[t+j]) = 0$ for $t > 0$, since $t+j-i \geq 2$. We also have that if $i > j$, then $\text{Hom}(T_i[i], T_j[j][t]) \simeq \text{Hom}(T_i, T_j[t-(i-j)])$. Hence, for $i > j$, we have that $\text{Hom}(T_i, T_j[t-(i-j)]) = 0$ for all $t > 0$ if and only if $\text{Hom}(T_i, T_j) = 0 = \text{Ext}^1(T_i, T_j)$. This finishes the proof. □

We now show the following connection between partial silting objects and exceptional sequences in $\text{mod } H$.

Lemma 2.2. (a) *Let T_0, \dots, T_m be partial tilting objects in $\text{mod } H$. As in (1), let $T = \bigoplus_{i=0}^m T_i[i]$, and let $\tilde{T} = \bigoplus_{i=0}^m T_i$. We have:*

- (i) *T is a partial silting object if and only if the summands of \tilde{T} can be ordered to be an exceptional sequence with all summands of T_i preceding all summands of T_j for $i < j$.*
- (ii) *T is a silting object if and only if the summands of \tilde{T} can be ordered as in (i) above to a complete exceptional sequence.*
- (b) *Any silting object has exactly n indecomposable direct summands, up to isomorphism.*
- (c) *Let $T' = T'_0 \oplus T'_1 \oplus \cdots \oplus T'_r$ be in $\text{mod } H$, with the T'_i indecomposable and mutually non-isomorphic, and let*

$$\overline{T'} = T'_0[0] \oplus T'_1[1] \oplus \cdots \oplus T'_r[r]$$

be in \mathcal{D} . Then we have that

- (i) $(T'_0, T'_1, \dots, T'_r)$ is an exceptional sequence if and only if $\overline{T'}$ is a partial silting object.
- (ii) $(T'_0, T'_1, \dots, T'_r)$ is a complete exceptional sequence if and only if $\overline{T'}$ is a silting object.

Proof. (a)(i) Assume T is partial tilting. Then each T_i is a partial tilting module over H , and hence the endomorphism algebras $\text{End}_H(T_i)$ has no oriented cycles [HR]. It now follows from Lemma 2.1 that the indecomposable direct summands of T_i can be ordered as $T_i^{(1)}, \dots, T_i^{(s)}$, with $\text{Hom}(T_i^{(u)}, T_i^{(v)}) = 0$ for $u > v$.

For the opposite implication, we assume there is an ordering such that $\text{Hom}(T_i^{(u)}, T_i^{(v)}) = 0$ for $u > v$. We then check the conditions of Lemma 2.1. Condition (i) holds by assumption. Condition (ii) follows from the definition of exceptional sequence.

(ii) Assume that T is a silting object, and that \tilde{T} can be ordered to an exceptional sequence \mathcal{E} . Assume \mathcal{E} is not complete. Then we can extend \mathcal{E} to a new exceptional sequence by adding an object U at the right end. Then $T \oplus U[r+1]$ is a partial silting object by part (i), and we have a contradiction, so \mathcal{E} must be complete.

On the other hand, assume that \tilde{T} can be ordered to an exceptional sequence \mathcal{E}_T , and that T is partial silting but not silting. Then there is some $R[j]$ in \mathcal{D} , with R not in $\text{add} T$, such that $T \oplus R[j]$ is partial silting. Note that j might be one of the shifts already occurring in the decomposition of T , in which case we add R to T_j . Then, using (i), we get a contradiction to the completeness of \mathcal{E}_T .

(b) This is a direct consequence of (a)(ii).

(c) Part (i) follows directly from Lemma 2.1, while part (ii) follows from (b) in combination with Lemma 2.1. □

2.2. Silting objects and maximal m -rigid objects. Here we show that silting objects are closely related to m -cluster tilting objects in m -cluster categories.

Recall that for an integer $m \geq 1$, the m -cluster category of a finite dimensional hereditary algebra H is the orbit category $\mathcal{C}_m = \mathcal{D}/\tau^{-1}[m]$, where τ is the AR-translation in \mathcal{D} , see [H], and $[m]$ is the m -fold composition of $[1]$, the shift functor. This category is known to be triangulated by [K]. It was first studied in the case $m = 1$ in [BMRRT]. An object T in \mathcal{C}_m is said to be *m -rigid* if $\text{Ext}_{\mathcal{C}_m}^i(T, T) = 0$ for $i = 1, \dots, m$; and if T is maximal with this property, then it is said to be *maximal m -rigid*. This has been shown [W, ZZ] to be equivalent to T being m -cluster tilting, that is $\text{Ext}_{\mathcal{C}_m}^i(T, X) = 0$ if and only if X is in $\text{add} T$. The maximal m -rigid objects are known to have exactly n non-isomorphic indecomposable summands [Z], see also [W].

Now \mathcal{C}_m is a Krull-Schmidt category, and we fix

$$\mathcal{S}_m = \text{mod } H[0] \vee \text{mod } H[1] \vee \dots \vee \text{mod } H[m-1] \vee H[m].$$

Then \mathcal{S}_m is a fundamental domain for \mathcal{C}_m in \mathcal{D} ; this means that the map from isomorphism classes of objects in \mathcal{S}_m to isomorphism classes of objects in \mathcal{C}_m is bijective.

It is easy to see the following (see [W, Lemma 1.1]).

Lemma 2.3. *Let T be an object in \mathcal{S}_m , then $\text{Ext}_{\mathcal{D}}^i(T, T) = 0$ for $i = 1, \dots, m$ if and only if $\text{Ext}_{\mathcal{C}_m}^i(T, T) = 0$ for $i = 1, \dots, m$.*

We then obtain the following.

Proposition 2.4. *Let H be a finite dimensional hereditary algebra, and let \mathcal{S}_m be the fundamental domain as above for the m -cluster category \mathcal{C}_m of H . Let T be an object in \mathcal{S}_m . Then we have the following:*

- (a) *T is a partial silting object in \mathcal{D} if and only if T is rigid in \mathcal{C}_m .*
- (b) *T is a silting object in \mathcal{D} if and only if T is an m -cluster tilting object in \mathcal{C}_m .*

Proof. For (a), assume first that T is a partial silting object, then it is m -rigid by Lemma 2.3.

Assume T is m -rigid, so that $\text{Ext}_{\mathcal{D}}^i(T, T) = 0$ for $i = 1, \dots, m$. Let X, Y be in $\text{mod } H$, and assume $X[r], Y[s]$ are in \mathcal{S}_m . If $r \neq m$, then $r - s < m$. So if $t > m$, then $t - (r - s) > 1$, and hence $\text{Ext}_{\mathcal{D}}^t(X[r], Y[s]) = \text{Hom}_{\mathcal{D}}(X, Y[t - (r - s)]) = 0$. If $r = m$, then for $t > m$ we have that $s + t - m \geq 1$, and we have $\text{Ext}_{\mathcal{D}}^t(X[m], Y[s]) = \text{Hom}_{\mathcal{D}}(X, Y[(s + t - m)])$. This vanishes, since it follows from $r = m$ that X must be projective.

(b): This follows from (a), since silting objects can be characterized among partial silting objects as those with n non-isomorphic indecomposable summands, and similarly for m -cluster tilting objects among m -rigid objects. \square

2.3. An example. The following example demonstrates that it is not possible to fix a placement of each exceptional H -module into \mathcal{D} sending exceptional sequences to silting objects.

Consider the following simple example. Let $H = kQ$, where Q is the quiver

$$1 \longleftarrow 2$$

Then $\text{mod } H$ has three indecomposable modules, the simples S_1 and S_2 , and the projective P_2 corresponding to vertex 2. The complete exceptional sequences are

$$(S_1, P_2), (P_2, S_2) \text{ and } (S_2, S_1)$$

Suppose the placements of S_1, P_2, S_2 are $S_1[a], P_2[b], S_2[c]$. It is straightforward to check the following:

- $S_1[a] \oplus P_2[b]$ is a silting object only if $a \leq b$
- $P_2[b] \oplus S_2[c]$ is a silting object only if $b \leq c$
- $S_2[c] \oplus S_1[a]$ is a silting object only if $c < a$

Clearly, we cannot have all three of the above conditions simultaneously. Hence, it is impossible to define a placement simultaneously for all exceptional H -modules taking exceptional sequences to silting objects.

3. CONNECTIONS BETWEEN MUTATIONS

In this section we compare the mutation operations in our three settings.

We have already discussed mutation for exceptional sequences. Recall that \mathcal{S}_m denotes the full subcategory of \mathcal{D} generated by

$$\text{mod } H[0] \vee \text{mod } H[1] \vee \dots \vee \text{mod } H[m-1] \vee H[m].$$

For an H -module X , we say that $X[v]$ has degree v , denoted by $d(X[v]) = v$. When we refer to the degree of an object in \mathcal{C}_m , we mean the degree of its lifting to \mathcal{S}_m . We say that a map $f: X \rightarrow Y$ in the m -cluster category is a D -map if it is induced by a map $X \rightarrow Y$ in the standard domain \mathcal{S}_m . Otherwise we say that it is an F -map.

The following is proved in [ZZ] (and (a) also in [W]).

Proposition 3.1. *Let $T = T_1 \oplus \cdots \oplus T_n$ be a cluster tilting object in \mathcal{C}_m . Fix j such that $1 \leq j \leq n$.*

- (a) *There are exactly $m+1$ non-isomorphic indecomposable objects M_0, \dots, M_m , such that $T/T_j \oplus M_i$ is an m -cluster tilting object.*
- (b) *The indecomposable objects M_0, \dots, M_m of part (a) can be numbered so that there are triangles*

$$M_{i-1} \xrightarrow{f_i} B_i \xrightarrow{g_i} M_i \rightarrow$$

for $i = 1, \dots, m$, with indices computed modulo m , such that each f_i is a minimal left $\text{add } T/T_j$ -approximation in \mathcal{C}_m and each g_i is a minimal right $\text{add } T/T_j$ -approximation in \mathcal{C}_m .

- (c) *For $0 \leq i \leq m-1$, we have that $d(M_i) \leq d(M_{i+1})$.*
- (d) *For $0 \leq i \leq m$, we have that $i-1 \leq d(M_i) \leq i$.*

By Proposition 2.4 it follows that an almost complete m -cluster tilting object T/T_j can be considered an almost complete silting object in \mathcal{D} , lying in \mathcal{S}_m . It also follows that the complements of T/T_j in \mathcal{C}_m coincide with those complements of T/T_j (as an almost complete silting object) which lie in \mathcal{S}_m .

In this section we show that mutation of m -cluster tilting objects can be naturally interpreted as a mutation of silting objects. We also point out some simple examples demonstrating that connections to mutation of exceptional sequences are more involved.

3.1. Mutation of silting objects and cluster tilting objects. We begin with a lemma, which is an adaption of a similar statement for tilting modules or cluster tilting objects. The proof is also similar.

Lemma 3.2. *Let \overline{T} be an almost complete silting object, and let M be a complement.*

- (a) *Let*

$$(2) \quad M^* \xrightarrow{f} B \xrightarrow{g} M \rightarrow$$

be a triangle such that g is a minimal right $\text{add } \overline{T}$ -approximation. Then M^ is a complement to \overline{T} , with $M^* \not\cong M$, and with f a minimal left $\text{add } \overline{T}$ -approximation.*

- (b) *Let*

$$(3) \quad M \xrightarrow{f'} B' \xrightarrow{g'} M^\# \rightarrow$$

be a triangle such that f' is a minimal left $\text{add } \overline{T}$ -approximation. Then $M^\#$ is a complement to \overline{T} , with $M^\# \not\cong M$, and with g' a minimal right $\text{add } \overline{T}$ -approximation.

Proof. We prove only (a); the proof of (b) is dual. By applying $\text{Hom}_{\mathcal{D}}(\overline{T}, _)$ and $\text{Hom}_{\mathcal{D}}(_, \overline{T})$ to the triangle (2), and considering the corresponding long exact sequences, we get that $\text{Ext}^i(\overline{T}, M^*) = \text{Ext}^i(M^*, \overline{T}) = 0$ for $i > 0$. Note that, in particular, we use that g is a right $\text{add } \overline{T}$ -approximation to obtain that $\text{Ext}^1(\overline{T}, M^*)$ vanishes.

From the exact sequence

$$\text{Hom}(B, \overline{T}) \rightarrow \text{Hom}(M^*, \overline{T}) \rightarrow \text{Ext}^1(M, \overline{T}) = 0$$

it follows that f is a left $\text{add } \overline{T}$ -approximation.

It is clear that f must be left minimal, since M is indecomposable, and not a direct summand in B .

We claim that M^* must be indecomposable. Assume to the contrary that $M^* = U \oplus V$, with U and V both non-zero. Consider minimal left $\text{add } \overline{T}$ -approximations $f_1: U \rightarrow B_1$ and $f_2: V \rightarrow B_2$, and complete to get triangles

$$U \rightarrow B_1 \rightarrow X \rightarrow \text{ and } V \rightarrow B_2 \rightarrow Y \rightarrow$$

Since the direct sum of the triangles is the triangle (2), it follows that $M \simeq X \oplus Y$, and hence either $X = 0$ or $Y = 0$. If $X = 0$, then $B_1 \rightarrow 0$ is a direct summand of $f: B \rightarrow M$, which contradicts that g is right minimal. Similarly $Y = 0$ leads to a contradiction, so we get that M^* is indecomposable.

Clearly M^* is not in $\text{add } \overline{T}$, since then f would be an isomorphism, contradicting $M \neq 0$.

We are now left with showing $\text{Ext}^i(M^*, M^*) = 0$ for $i > 0$. Applying $\text{Hom}(_, M)$ to the triangle (2), considering the corresponding long exact sequence, and using that $\text{Ext}^1(M, M) = 0$, we get an epimorphism

$$(4) \quad \text{Hom}(B, M) \rightarrow \text{Hom}(M^*, M)$$

So any map $h: M^* \rightarrow M$ factors through $f: M^* \rightarrow B$.

Then we apply $\text{Hom}(M^*, _)$ to the triangle (2) and consider the resulting exact sequence

$$\text{Hom}(M^*, B) \rightarrow \text{Hom}(M^*, M) \rightarrow \text{Ext}^1(M^*, M^*) \rightarrow \text{Ext}^1(M^*, B)$$

Since the rightmost term vanishes, it is sufficient to show that $\text{Hom}(M^*, B) \rightarrow \text{Hom}(M^*, M)$ is surjective in order to obtain $\text{Ext}^1(M^*, M^*) = 0$. To see that any map $h: M^* \rightarrow M$ factors through $g: B \rightarrow M$, consider the commutative diagram

$$\begin{array}{ccccc} & & & & B \\ & & & & \downarrow t \\ & & & \swarrow s & \\ M^* & \xrightarrow{f} & B & \xrightarrow{g} & M \\ & \downarrow h & \swarrow t & & \\ & M & & & \end{array}$$

Here t is obtained from the epimorphism (4), and we get $s: B \rightarrow B$ by using that g is a right $\text{add } \overline{T}$ -approximation. So $h = tf = gsf$.

Applying $\text{Hom}(_, M)$ to the triangle (2), and considering the resulting long exact sequence, we obtain $\text{Ext}^i(M^*, M) = 0$ for $i \geq 1$.

It then follows from the long exact sequence obtained by applying $\text{Hom}(M^*, _)$ to the triangle (2) that $\text{Ext}^i(M^*, M^*) = 0$ for $i > 1$. This finishes the proof. \square

The triangles appearing in Lemma 3.2 are called *exchange triangles* in \mathcal{D} . We need the following observation.

Lemma 3.3. *If a minimal left $\text{add } \overline{T}$ -approximation α in \mathcal{C}_m is a D -map, then it is also a minimal left $\text{add } \overline{T}$ -approximation in \mathcal{D} .*

For our main result in this section, we also need the following two lemmas.

Lemma 3.4. *Let \overline{T} be an almost complete sifting object with a complement M . Let $m \geq 2$, and let*

$$(5) \quad M^* \xrightarrow{f} B \xrightarrow{g} M \rightarrow$$

be an exchange triangle in \mathcal{C}_m with $B \neq 0$. Then $d(M^) \leq d(M)$ if and only if both f and g are D -maps.*

Proof. If both f and g are D -maps, then clearly $d(M^*) \leq d(M)$.

Note that if $X \rightarrow Y$ is not a D -map, with X and Y indecomposable, we must have $d(Y) = 0$ and $d(X) \geq m - 1$.

Assume f is not a D -map. Then the middle term B must have a summand of degree 0, and hence $d(M) \leq 1$. Also, we have that $d(M^*)$ must be $m - 1$ or m . If $m \geq 3$, then clearly $d(M^*) > d(M)$. Consider the case $m = 2$. If $d(M^*) = 2$, then trivially $d(M^*) > d(M)$. Assume $d(M^*) = 1$. We already know that $d(M) \neq 2$, and now we claim that $d(M) \neq 1$. Assume $d(M) = 1$. The exchange triangle (5) must be induced either (i) by an exact sequence in $\text{mod } H$, or (ii) by a triangle

$$M^* \xrightarrow{f} B' \xrightarrow{g} \tau^{-1}M[2] \rightarrow$$

in the derived category \mathcal{D} . Then (i) is not possible, by the assumption that f is an F -map. Also (ii) is not possible, since a map $\tau^{-1}M[2] \rightarrow M^*[1]$ is zero in the derived category, and hence the triangle must split, which is a contradiction. Hence $d(M) = 0$.

Assume f is a D -map, but g is not. Since g is not a D -map, then B must have a direct summand in degree $m - 1$ or m , and $d(M) = 0$. Since f is D -map we have $d(M^*) \geq m - 2$. So for $m \geq 3$, we clearly have $d(M^*) > d(M)$. Consider the case $m = 2$, and assume $d(M^*) = d(M) = 0$. Then B must have a summand of degree > 0 , and as above this implies that the exchange triangle (5) must be a triangle

$$M^* \xrightarrow{f} B' \xrightarrow{g} \tau^{-1}M[2] \rightarrow$$

in the derived category \mathcal{D} , which is not possible, since a map $\tau^{-1}M[2] \rightarrow M^*[1]$ is zero in the derived category, and hence the triangle must split, which is a contradiction. \square

We can now prove the main result in this section.

Theorem 3.5. *Let $T = T_1 \oplus \cdots \oplus T_n$ be a basic siltng object in $\mathcal{D} = D^b(H)$, where the T_i are indecomposable and n is the number of isomorphism-classes of simple H -modules. We assume without loss of generality that T is in \mathcal{D}^+ . Let m be an integer such that T is in the fundamental domain \mathcal{S}_m of the m -cluster category \mathcal{C}_m . Fix an indecomposable direct summand T_i , with $1 \leq i \leq n$.*

- (a) *The almost complete siltng object T/T_i has exactly $m + 1$ non-isomorphic complements M_0, M_1, \dots, M_m lying in \mathcal{S}_m , which are ordered such that $d(M_i) \leq d(M_{i+1})$.*

- (b) *For each $j = 1, \dots, m$, there is a triangle in \mathcal{D}*

$$(6) \quad M_{j-1} \xrightarrow{f_j} B_j \xrightarrow{g_j} M_j \rightarrow,$$

such that f_j is a minimal left add T/T_i -approximation, and g_j is a minimal right add T/T_i -approximation.

- (c) *The triangles (6) are also exchange triangles in \mathcal{C}_m .*

Note that in the m -cluster category \mathcal{C}_m , there is an additional exchange triangle

$$M_m \rightarrow B_0 \rightarrow M_0 \rightarrow$$

which is not a triangle in \mathcal{D} .

We have the following direct consequence, with assumptions as in the theorem.

Corollary 3.6. *T/T_i has a countably infinite number of non-isomorphic complements M_i for $i \in \mathbb{Z}$. In particular, there are complements M_{-1} and M_{m+1} , such that $M_j \simeq M_{-1}[j + 1]$ for $j < -1$, and $M_j = M_{m+1}[j - (m + 1)]$ for $j > m + 1$.*

Proof of theorem. Note that by Proposition 2.4, the complements of T/T_i as an almost complete siltng object and as an almost complete m -cluster tilting object coincide. From Proposition 3.1, there are, up to isomorphism, exactly $m + 1$ complements M_0, M_1, \dots, M_m of T/T_i as an almost complete m -cluster tilting object, and they can be ordered such that for $j = 0, \dots, m - 1$, there are exchange triangles $M_j \rightarrow B_{j+1} \rightarrow M_{j+1} \rightarrow$ in \mathcal{C}_m and $d(M_j) \leq d(M_{j+1})$.

Assume first $m \geq 2$. All the maps $M_j \rightarrow B_{j+1}$ and $B_{j+1} \rightarrow M_{j+1}$ are then \overline{D} -maps by Lemma 3.4. Then $M_0 \rightarrow B_1$ is by Lemma 3.3 a minimal left add \overline{T} approximation in \mathcal{D} . Hence, by Lemma 3.2 (b) it follows that the exchange triangle $M_0 \rightarrow B_1 \rightarrow M_1$ in \mathcal{D} coincides with the exchange triangle in \mathcal{C}_m .

Iterating this for $M_i \rightarrow B_{i+1}$, all of (a), (b), and (c) follow.

Consider now the case $m = 1$, and the exchange sequence

$$(7) \quad M_0 \rightarrow B_1 \rightarrow M_1 \rightarrow$$

in \mathcal{C}_m .

(i) If $d(M_0) = d(M_1) = 0$ (up to reordering M_0 and M_1) the sequence (7) is induced by an exact sequence in $\text{mod } H$.

(ii) If $d(M_0) = 0$ and $d(M_1) = 1$, it is straightforward to check that the sequence (7) must be induced by a sequence $M_0 \rightarrow B_0 \rightarrow P[1] \rightarrow$ in the derived category, with both $M_0 \rightarrow B_1$ and $B_1 \rightarrow P[1]$ being D -maps, and with P an indecomposable projective.


Hence in both case (i) and (ii) both (a), (b) and (c) follow from combining Lemmas 3.2 and 3.3 as in the above case. \square

3.2. Example. Let $\mathcal{E} = \mathcal{E}_0$ be a complete exceptional sequence and let $\mathcal{E}_i = \mu_i(\mathcal{E})$ for $i = 1, \dots, n-1$. A natural question is: Is it possible to embed these n exceptional sequences simultaneously in \mathcal{S}_m for some value of m , such that the following are satisfied

- each X occurring in some \mathcal{E}_i is mapped to a fixed $X[j_X]$
- each \mathcal{E}_i is mapped to an m -cluster tilting object U_i , in such a way that
- the m -cluster tilting object U_i is obtained by mutation of U_0 in \mathcal{C}_m

The following example demonstrates that such an embedding is not possible in general.

Let Q be the quiver



and $H = kQ$. Let P_i be the indecomposable projective corresponding to vertex i . Then $\mathcal{E} = (P_1, P_2, P_3)$ is a complete exceptional sequence. We have $\mu_1(\mathcal{E}) = (P_2, S_2, P_3)$ and $\mu_2(\mathcal{E}) = (P_1, P_3, R)$, where S_1 is the simple corresponding to vertex 1, and R is a regular indecomposable module with composition factors S_3 and S_1 .

Assume the images of the P_i are $P_1[a], P_2[b], P_3[c]$. We must have that the image $S_2[d]$ of S_2 is obtained by mutation of the cluster tilting object $P_1[a] \oplus P_2[b] \oplus P_3[c]$ at $P_1[a]$. Hence we must have $\text{Hom}(P_1[a], P_2[b]) \neq 0$ and $\text{Hom}(P_1[a], P_3[c]) = 0$, so we must have $a = b$ and $a \neq c$. Considering mutation at $P_2[b]$ we get similarly that $\text{Hom}(P_2[b], P_2[c]) \neq 0$, so we must have $b = c$, and we have a contradiction.

4. PLACEMENT OF AN ALMOST COMPLETE EXCEPTIONAL SEQUENCE AND ITS COMPLEMENTS

As before, let H be a hereditary algebra with n isomorphism classes of simple modules. In this section we show that for a fixed almost complete exceptional sequence \mathcal{E} in $\text{mod } H$, and its set of n complements, there is a natural interpretation of \mathcal{E} as an almost complete silting object \widehat{A} in \mathcal{S}_{n-1} , in such a way that the complements and exchange sequences correspond.

Consider an almost complete exceptional sequence $\mathcal{E} = (A_0, A_1, \dots, A_{n-2})$. Let $\mathcal{C} = \{C_0, C_1, \dots, C_{n-1}\}$ be the complements of \mathcal{E} , such that each

$$\mathcal{E}_i = (A_0, A_1, \dots, C_i, A_i, \dots, A_{n-2})$$

is a complete exceptional sequence. Recall from section 1.1 that for each $i = 0, \dots, n-2$, there is a mutation μ_i , replacing (C_i, A_i) with (A_i, C_{i+1}) and that there is an induced triangle in \mathcal{D}

$$(8) \quad C_i \xrightarrow{f_i} A_i^{r_i}[v] \xrightarrow{g_i} C_{i+1}[w] \rightarrow,$$

with f_i a minimal left (thick A_i)-approximation, with $v, w \in \{0, 1\}$. Note that f_i might be the zero-map (i.e. $r_i = 0$), in which case $C_{i+1} = C_i$ and $w = 1$.

We want to define objects $\widehat{C}_i, \widehat{A}_i$ inside the standard domain \mathcal{S}_{n-1} , such that the following hold.

- $\widehat{C}_i = C_i[t_i]$ and $\widehat{A}_i = A_i[u_i]$, for integers t_i, u_i such that $\widehat{C}_i, \widehat{A}_i$ in \mathcal{S}_{n-1} .
- $\bigoplus_{i=0}^{n-2} \widehat{A}_i$ is an almost complete silting object with complements \widehat{C}_i .
- There are exchange triangles $\widehat{C}_i \rightarrow \widehat{A}_i^{r_i} \rightarrow \widehat{C}_{i+1} \rightarrow$ given by shifting the mutation triangles (8).

Theorem 4.1. *Given an almost complete exceptional sequence $\mathcal{E} = (A_0, A_1, \dots, A_{n-2})$, there exists a placement of the objects A_i, C_i in \mathcal{S}_m with the above properties.*

Proof. The proof is by induction on n . In case $n = 1$, the almost complete exceptional sequence \mathcal{E} is empty, and has a unique complement $C_0 = H$. The standard domain \mathcal{S}_0 also consists of only the object H , and a placement trivially exists.

Assume now $n > 1$, and consider the category

$$A_{n-2}^\perp = \{U \in \text{mod } H \mid \text{Hom}(A_{n-2}, U) = \text{Ext}^1(A_{n-2}, U) = 0\}.$$

By [CB, R] there is an hereditary algebra H' such that $\text{mod } H' \simeq A_{n-2}^\perp$, and H' has $n - 1$ isomorphism classes of simple modules.

Since each A_i for $0 \leq i \leq n - 3$ and each C_i for $0 \leq i \leq n - 2$ is in A_{n-2}^\perp , we have by induction a placement for these in the fundamental domain for H' , which as a full additive subcategory of \mathcal{D} is generated by $A_{n-2}^\perp[j]$ for $0 \leq j \leq n - 3$, and $X[n - 2]$, where X is the direct sum of all indecomposable Ext-projectives of A_{n-2}^\perp .

We now define \widehat{A}_{n-2} and \widehat{C}_{n-1} according to the following rules.

- (P1) If $\text{Hom}(C_{n-2}, A_{n-2}) \neq 0$, then let $d(\widehat{A}_{n-2}) = d(\widehat{C}_{n-2})$.
- (P2) If $\text{Ext}^1(C_{n-2}, A_{n-2}) \neq 0$, then let $d(\widehat{A}_{n-2}) = d(\widehat{C}_{n-2}) + 1$.
- (P3) Otherwise let $d(\widehat{A}_{n-2}) \geq n - 3$ and minimal such that the following is satisfied
 - (i) $d(\widehat{A}_{n-2}) \geq d(\widehat{C}_{n-2})$,
 - (ii) $d(\widehat{A}_{n-2}) > d(\widehat{A}_{n-3})$ if $\text{Ext}^1(A_{n-3}, A_{n-2}) \neq 0$,
 - (iii) $d(\widehat{A}_{n-2}) > d(\widehat{A}_{n-4})$ if $\text{Ext}^1(A_{n-4}, A_{n-2}) \neq 0$.

In case (P1) or (P2), \widehat{C}_{n-1} is defined such that we have a triangle

$$\widehat{C}_{n-2} \rightarrow \widehat{A}_{n-2}^{r_{n-2}} \rightarrow \widehat{C}_{n-1} \rightarrow .$$

In case (P3), we let $\widehat{C}_{n-1} = \widehat{C}_{n-2}[1]$.

We claim that with this definition of \widehat{A}_{n-2} and \widehat{C}_{n-1} we have that the following hold.

- \mathcal{U}_n : \widehat{A}_{n-2} is in \mathcal{S}_{n-1} .
- \mathcal{V}_n : $\widehat{A} = \bigoplus_{i=0}^{n-2} \widehat{A}_i$ is an almost complete silting object.
- \mathcal{W}_n : \widehat{C}_i for $i = 0, \dots, n - 1$ are the complements of \widehat{A} inside \mathcal{S}_{n-1} .
- \mathcal{X}_n : If $d(\widehat{A}_{n-2}) = d(\widehat{C}_{n-1})$, then there is a directed path from A_{n-2} to C_{n-1} in the Hom-Ext quiver \mathcal{Q} associated to the exceptional sequence

$$(A_0, \dots, A_{n-2}, C_{n-1})$$

with the property that the last arrow on this path corresponds to a morphism to C_{n-1} .

$$\mathcal{Y}_n: d(\widehat{C_{n-1}}) \geq d(\widehat{A_{n-2}}) \geq d(\widehat{C_{n-2}})$$

It is clear that all statements hold for $n = 1$. It is also clear that the claim of the theorem follows from $\mathcal{U}_n, \mathcal{V}_n, \mathcal{W}_n$. We need also to include \mathcal{X}_n and \mathcal{Y}_n as part of our induction argument.

We first need a lemma. In [B] it was shown that for any exceptional non-projective module Y , there is a (Bongartz-)complement W , with the properties that

- $Y \oplus W$ is tilting,
- $\text{Ext}^1(Y, U) = 0$ implies $\text{Ext}^1(W, U) = 0$ for any U in $\text{mod } H$ and
- $\text{Hom}(Y, W) = 0$.

Lemma 4.2. *Assume A_{n-2} is not projective, then the direct sum X of one copy of each of the indecomposable Ext-projectives in A_{n-2}^\perp , up to isomorphism, equals the Bongartz-complement of A_{n-2} .*

Proof. Let W be the Bongartz complement of A_{n-2} in $\text{mod } H$. Then it follows directly from the properties of the Bongartz complement that W is in A_{n-2}^\perp , and is Ext-projective in this subcategory. Since it is known that A_{n-2}^\perp is equivalent to a module category with $n-1$ isomorphism classes of simples, the statement follows. \square

By induction we can assume $\mathcal{U}_m\text{-}\mathcal{Y}_m$ hold for $m < n$. We need to prove that they hold also for n .

Proof of \mathcal{U}_n :

It is clear that we only need to consider the case where A_{n-2} is placed according to P2 or P3. Assume first $\text{Ext}^1(C_{n-2}, A_{n-2}) \neq 0$, so that $d(\widehat{A_{n-2}}) = d(\widehat{C_{n-2}}) + 1$ according to P2. If $d(\widehat{C_{n-2}}) \leq n-3$, then $d(\widehat{A_{n-2}}) \leq n-2$. So assume $d(\widehat{C_{n-2}}) = n-2$ and $d(\widehat{A_{n-2}}) = n-1$. If A_{n-2} is non-projective, then C_{n-2} must be a summand in X , and hence Ext-projective in A_{n-2}^\perp . Hence, by Lemma 4.2 we have that $\text{Ext}^1(C_{n-2}, A_{n-2}) = 0$, contradicting our assumption. If A_{n-2} is projective, then $\widehat{A_{n-2}}$ is in \mathcal{S}_n , and we are done.

Assume $\widehat{A_{n-2}}$ is placed according to case P3. We need to show that if $d(\widehat{A_{n-2}}) = n-1$, then A_{n-2} is projective. We have that $d(\widehat{C_{n-2}}) \leq n-2$, by \mathcal{W}_{n-1} . Therefore, if $d(\widehat{A_{n-2}}) = n-1$, this must be forced by (ii) or (iii) in P3. However $d(\widehat{A_{n-4}}) \leq n-3$ by \mathcal{U}_{n-2} , so we need only consider option (ii), i.e. $\text{Ext}^1(A_{n-3}, A_{n-2}) \neq 0$. We must have $d(\widehat{A_{n-3}}) = n-2$, hence A_{n-3} is a summand in X , the sum of the Ext-projectives in A_{n-2}^\perp . If A_{n-2} was non-projective, then $\text{Ext}^1(A_{n-3}, A_{n-2}) = 0$ by Lemma 4.2, and we have a contradiction. Hence, we have that A_{n-2} is projective, and therefore $\widehat{A_{n-2}}$ is in \mathcal{S}_{n-1} , and we are done.

Proof of \mathcal{V}_n :

By induction $\widehat{A'} = \bigoplus_{i=0}^{n-3} \widehat{A}_i$ is a partial silting object in $\mathcal{D}(H')$, lying inside the fundamental domain for H' . Clearly, $\widehat{A'}$ is also a partial silting object in $\mathcal{D}(H)$.

We need to show that if $d(\widehat{A_{n-2}}) = d(\widehat{A_{n-j}})$ for some $j \geq 3$, then $\text{Ext}^1(A_{n-j}, A_{n-2}) = 0$. Note that for $j \geq 5$, we have that $d(\widehat{A_{n-j}}) \leq n-4$ by \mathcal{U}_{n-3} , while $d(\widehat{A_{n-2}}) \geq n-3$, so we only need to consider the cases $j = 3, 4$.

We first consider the case $j = 3$. Assume $d(\widehat{A_{n-2}}) = d(\widehat{A_{n-3}})$. By \mathcal{Y}_{n-1} , we have $d(\widehat{C_{n-2}}) \geq d(\widehat{A_{n-3}})$, and by considering the different cases of our placement algorithm $d(\widehat{A_{n-2}}) \geq d(\widehat{C_{n-2}})$. Hence we must have $d(\widehat{A_{n-2}}) = d(\widehat{C_{n-2}}) = d(\widehat{A_{n-3}})$.

Using \mathcal{X}_{n-1} , we find, in the Hom-Ext quiver \mathcal{Q} associated to the exceptional sequence $(A_0, \dots, A_{n-3}, C_{n-2})$ in $\text{mod } H'$, a directed path from A_{n-3} to C_{n-2} . Assume first we are in case P1, so that we have a non-zero map from C_{n-2} to A_{n-2} . If $\text{Ext}^1(A_{n-3}, A_{n-2}) \neq 0$, this would give us an arrow from A_{n-2} to A_{n-3} in the Hom-Ext quiver \mathcal{Q} , and hence an oriented cycle. By Theorem 1.4, such cycles do not exist, so we must have $\text{Ext}^1(A_{n-3}, A_{n-2}) = 0$, which is what we wanted to show. Now assume we are in case P3. Then, since $d(\widehat{A_{n-2}}) = d(\widehat{A_{n-3}})$, we must have that $\text{Ext}^1(A_{n-3}, A_{n-2}) = 0$, according to (ii). It is also clear we cannot be in case P2, since $d(\widehat{A_{n-2}}) = d(\widehat{C_{n-2}})$.

We now consider the case $j = 4$. Assume $d(\widehat{A_{n-2}}) = d(\widehat{A_{n-4}})$. Using \mathcal{Y}_{n-1} and \mathcal{Y}_{n-2} , this implies that

$$d(\widehat{A_{n-2}}) = d(\widehat{C_{n-2}}) = d(\widehat{A_{n-3}}) = d(\widehat{C_{n-3}}) = d(\widehat{A_{n-4}}).$$

We can apply \mathcal{X}_{n-2} to find a directed path from A_{n-4} to C_{n-3} in the Hom-Ext quiver \mathcal{Q} of

$$(A_0, \dots, A_{n-4}, C_{n-3}).$$

Since $d(\widehat{C_{n-2}}) = d(\widehat{C_{n-3}})$, there must be an exchange triangle

$$\widehat{C_{n-3}} \rightarrow \widehat{A_{n-3}^{r_{n-3}}} \rightarrow \widehat{C_{n-2}} \rightarrow$$

which is induced by an exact sequence in $\text{mod } H$. Since C_{n-3} embeds into $A_{n-3}^{r_{n-3}}$, we can compose the final morphism in the path from A_{n-4} to C_{n-3} with this inclusion, obtaining a path from A_{n-4} to A_{n-3} , onto which the arrow from A_{n-3} to C_{n-2} can be added, giving us a path from A_{n-4} to C_{n-2} .

Assume we are in case P1. We then have a map from C_{n-2} to A_{n-2} . Hence, if $\text{Ext}^1(A_{n-4}, A_{n-2}) \neq 0$, this would give us an arrow from A_{n-2} to A_{n-4} , and hence an oriented cycle in the Hom-Ext quiver \mathcal{Q} . Therefore we must have $\text{Ext}^1(A_{n-4}, A_{n-2}) = 0$. Assume we are in case P3. Then, since $d(\widehat{A_{n-2}}) = d(\widehat{A_{n-4}})$, we have that $\text{Ext}^1(A_{n-4}, A_{n-2}) = 0$, according to (iii). It is also clear we cannot be in case P2, since $d(\widehat{A_{n-2}}) = d(\widehat{C_{n-2}})$. Hence we are done with \mathcal{V}_n .

Proof of \mathcal{W}_n :

We first claim that $\widehat{C_{n-2}}$ is a complement to \widehat{A} . By induction, we only need to show that it is compatible with $\widehat{A_{n-2}}$. But this follows directly from P2 in our placement algorithm.

Next, we claim that all \widehat{C}_i for $i = 0, \dots, n-3$ are also complements. For this, consider the triangles

$$\widehat{C}_{i-1} \rightarrow \widehat{A}_i^{r_i} \rightarrow \widehat{C}_i \rightarrow$$

for $i = 1, \dots, n-2$. The maps $\widehat{A}_i^{r_i} \rightarrow \widehat{C}_i$ are by assumption minimal right \widehat{A}' -approximations in the fundamental domain for A_{n-2}^\perp , which is a full subcategory of \mathcal{D} . Hence they are also minimal right \widehat{A} -approximations in \mathcal{D} . We have that $\text{Hom}(\widehat{A}_{n-2}, \widehat{C}_j) = \text{Ext}^1(\widehat{A}_{n-2}, \widehat{C}_j) = 0$ for $j \leq n-2$ by our assumptions. Hence the maps $\widehat{A}_i \rightarrow \widehat{C}_i$ are also right \widehat{A} -approximations. Using Lemma 3.2, we obtain that all \widehat{C}_i for $i \leq n-2$ are complements.

Finally, we claim that \widehat{C}_{n-1} is also a complement. This follows directly from our placement algorithm, since the left \widehat{A}_{n-2} -approximation of \widehat{C}_{n-2} is indeed a left \widehat{A} -approximation. This follows since \widehat{C}_{n-2} does not have any non-zero maps to \widehat{A}' .

Proof of \mathcal{X}_n :

Assume first $\text{Hom}(C_{n-2}, A_{n-2}[j]) \neq 0$ for $j = 0$ or $j = 1$. Then $\widehat{A}_{n-2}, \widehat{C}_{n-1}$ are placed using P1 or P2, so if $d(\widehat{A}_{n-2}) = d(\widehat{C}_{n-1})$, then clearly there is a map $A_{n-2} \rightarrow C_{n-1}$.

So we can assume $\text{Hom}(C_{n-2}, A_{n-2}[j]) = 0$ for $j = 0, 1$, and hence $C_{n-1} = C_{n-2}$. Using \mathcal{W}_{n-1} and Proposition 3.1(d), we have that $d(\widehat{C}_{n-2}) \geq n-3$, and hence $d(\widehat{C}_{n-1}) \geq n-2$. Hence $d(\widehat{A}_{n-2}) \geq n-2$ by assumption. Since $d(\widehat{A}_{n-2}) \neq n-3$, one of the following must hold

- (i') $d(\widehat{C}_{n-2}) = d(\widehat{A}_{n-2}) = n-2$
- (ii') $\text{Ext}^1(A_{n-3}, A_{n-2}) \neq 0$ and (ii) in P3 applies
- (iii') $\text{Ext}^1(A_{n-4}, A_{n-2}) \neq 0$ and (iii) in P3 applies

Assume first (i'). In this case we have $d(\widehat{A}_{n-2}) = d(\widehat{C}_{n-2}) = n-2$ and $d(\widehat{C}_{n-2}) < d(\widehat{C}_{n-1})$, hence we cannot have $d(\widehat{A}_{n-2}) = d(\widehat{C}_{n-1})$, so there is nothing to show.

Assume now (ii'). By assumption $\widehat{C}_{n-1} = \widehat{C}_{n-2}[1]$, so we have that $d(\widehat{C}_{n-2}) = d(\widehat{A}_{n-3})$. Hence, we can apply \mathcal{X}_{n-1} , to get a directed path from A_{n-3} to C_{n-2} in the Hom-Ext quiver of the exceptional sequence

$$(A_0, \dots, A_{n-3}, C_{n-2})$$

and such that this path ends with an arrow corresponding to a homomorphism. Since $C_{n-2} = C_{n-1}$, this gives us a directed path from A_{n-3} to C_{n-1} in the Hom-Ext quiver of the exceptional sequence

$$(A_0, \dots, A_{n-3}, A_{n-2}, C_{n-1}).$$

By assumption, $\text{Ext}^1(A_{n-3}, A_{n-2}) \neq 0$, so we can add $A_{n-2} \rightarrow A_{n-3}$ to the front of the directed path, producing the directed path we need.

Assume now (iii'). By \mathcal{U}_{n-2} , we have $d(\widehat{A}_{n-4}) \leq n-3$. Then, since (iii) forced $d(\widehat{A}_{n-2}) > n-3$, we must have $d(\widehat{A}_{n-4}) = n-3$, and we have $d(\widehat{A}_{n-2}) = n-2$. By assumption, then $d(\widehat{C}_{n-1}) = n-2$, and since $C_{n-2} = C_{n-1}$, we have $d(\widehat{C}_{n-2}) = n-3$.

By \mathcal{Y}_{n-1} , we then have that also $d(\widehat{C_{n-3}}) = n - 3$. Since $d(\widehat{C_{n-2}}) = d(\widehat{C_{n-3}})$, we must have a short exact sequence

$$0 \rightarrow C_{n-3} \rightarrow A_{n-3}^{r_{n-3}} \rightarrow C_{n-2} \rightarrow 0$$

By \mathcal{X}_{n-2} we have a path ρ from A_{n-4} to C_{n-3} in the Hom-Ext quiver of the exceptional sequence

$$(A_0, \dots, A_{n-4}, C_{n-3})$$

ending in an arrow, say $A_j \rightarrow C_{n-3}$ for some j , and where this arrow is either an m-arrow or an e-arrow (i.e., it represents a morphism from A_j). We can prepend to ρ the x-arrow from A_{n-2} to A_{n-4} , resulting in a directed path from A_{n-2} to C_{n-3} .

We know that C_{n-3} injects into $A_{n-3}^{r_{n-3}}$, so we can compose the map from A_j to C_{n-3} with this inclusion, to obtain an arrow in the Hom-Ext quiver from A_j to A_{n-3} . There is also an arrow from A_{n-3} to C_{n-2} . Thus we can remove the last step of ρ and replace it by $A_j \rightarrow A_{n-3} \rightarrow C_{n-2} = C_{n-1}$, obtaining a directed path as desired.

Proof of \mathcal{Y}_n :

The inequality $d(\widehat{A_{n-2}}) \geq d(\widehat{C_{n-2}})$ follows directly from (P1), (P2) and part(i) of (P3).

The inequality $d(\widehat{C_{n-1}}) \geq d(\widehat{A_{n-2}})$ is clearly satisfied if we are in case (P1) or (P2). So assume we are in case (P3), and hence $C_{n-2} = C_{n-1}$, and $d(\widehat{C_{n-2}}) = d(\widehat{C_{n-1}}) - 1$.

Proposition 3.1 and \mathcal{W}_{n-1} together imply that $d(\widehat{C_{n-2}}) \geq n - 3$. Hence, we have $d(\widehat{A_{n-2}}) \geq n - 3$. Assume that the desired inequality is not satisfied, so we have $d(\widehat{C_{n-1}}) < d(\widehat{A_{n-2}})$. Then we must have $d(\widehat{A_{n-3}}) = d(\widehat{C_{n-1}}) > d(\widehat{C_{n-2}})$, which contradicts \mathcal{Y}_{n-1} . This finishes the proof for \mathcal{Y}_n . □

REFERENCES

- [AST] I. Assem, M.J. Souto Salorio and S. Trepode, *Ext-projectives in suspended subcategories*, Journal of pure and applied algebra **212** (2008), 423–434
- [B] K. Bongartz, *Tilted algebras*, Representations of algebras (Puebla, 1980), pp. 26–38, Lecture Notes in Math., 903, Springer, Berlin-New York, 1981
- [CB] W. Crawley-Boevey, *Exceptional sequences of representations of quivers*, Representations of algebras (Ottawa, ON, 1992), 117–124, CMS Conf. Proc., 14, Amer. Math. Soc., Providence, RI, 1993
- [BMRRT] A. B. Buan, R. J. Marsh, M. Reineke, I. Reiten and G. Todorov, *Tilting theory and cluster combinatorics*, Advances in Mathematics **204** (2) (2006), 572–618
- [H] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988. x+208 pp. ISBN: 0-521-33922-7
- [HR] D. Happel and C. M. Ringel, *Tilted algebras*, Trans. Amer. Math. Soc. 274 (1982), no. 2, 399–443
- [K] B. Keller, *On triangulated orbit categories*, Doc. Math. 10 (2005), 551–581
- [KV] B. Keller and D. Vossieck, *Aisles in derived categories*, Bull. Soc. Math. Belg. Ser. A 40 (1988), no. 2, 239–253

- [RS] C. Riedtmann and A. Schofield, *On open orbits and their complements*, J. Algebra 130 (1990), no. 2, 388–411
- [R] C. M. Ringel, *The braid group action on the set of exceptional sequences of a hereditary Artin algebra*, Abelian group theory and related topics (Oberwolfach, 1993), 339–352, Contemp. Math., 171, Amer. Math. Soc., Providence, RI, 1994
- [Ru] A. Rudakov, *Exceptional collections, mutations and helices. Helices and vector bundles*, 1–6, London Math. Soc. Lecture Note Ser., 148, Cambridge Univ. Press, Cambridge, 1990
- [T] H. Thomas, *Defining an m -cluster category*, J. Algebra 318 (2007), no. 1, 37–46
- [W] A. Wraalsen, *Rigid objects in higher cluster categories*, J. Algebra 321 (2009), no. 2, 532–547
- [Z] B. Zhu, *Generalized cluster complexes via quiver representations*, J. Algebraic Combin. 27 (2008), no. 1, 35–54
- [ZZ] Y. Zhou and B. Zhu, *Cluster combinatorics of d -cluster categories*, J. Algebra 321 (2009), no. 10, 2898–2915

INSTITUTT FOR MATEMATISKE FAG, NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET, N-7491 TRONDHEIM, NORWAY

E-mail address: `aslakb@math.ntnu.no`

INSTITUTT FOR MATEMATISKE FAG, NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET, N-7491 TRONDHEIM, NORWAY

E-mail address: `idunr@math.ntnu.no`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW BRUNSWICK, FREDERICTON NB, E3B 1J4 CANADA

E-mail address: `hthomas@unb.ca`